

Procedure Information Sheet – Laparoscopic Tubal Occlusion

Hosp No. :	HKID No.:
Case No. :	
Name :	
DOB :	M / F
Adm Date :	
Contact No. :	

Clinical diagnosis: unwanted fertility

Indication: unwanted fertility

1. Nature of the operation

- 1.1. General anaesthesia.
- 1.2. Pneumoperitoneum created by insufflation of carbon dioxide.
- 1.3. Incisions made.
- 1.4. Telescope and instruments passed into abdomen.
- 1.5. Local anaesthetic applied to fallopian tubes.
- 1.6. Fallopian tubes interrupted with Falope rings / clips.
- 1.7. Incisions closed.
- 1.8. All tissue removed will be sent to the Department of Pathology or disposed of as appropriate unless otherwise specified.
- 1.9. Photographic and/or video images may be recorded during the operation for education/research purpose. Please inform our staff if you have any objection.

2. Benefits of the procedure

- 2.1. Effective contraception.

3. Other consequences (after the procedure)

- 3.1. No effect on hormonal status in the presence of normal ovaries.
- 3.2. Coitus is not affected.
- 3.3. Irreversible method of contraception unless tubal reanastomosis done.
- 3.4. (May have unrelated menstruation change).

4. Risks and complications (may include, but are not limited to the following)

- 4.1. Women who are obese, who have significant pathology, who have undergone previous surgery or who have pre-existing medical conditions must understand that the quoted risks for serious or frequent complications will be increased.
- 4.2. Anaesthetic complications
- 4.3. Serious
 - 4.3.1. Failure to gain entry into the abdomen and to complete the intended procedure, requiring laparotomy
 - 4.3.2. Bleeding, may require blood transfusion
 - 4.3.3. Uterine perforation
 - 4.3.4. Injuries to the bowel, bladder or blood vessels (3 in every 1000, uncommon)
 - 4.3.5. One in every 12000 woman undergoing laparoscopy dies as a result of complications
 - 4.3.6. Pelvic infection
 - 4.3.7. Luteal phase pregnancy (2-3 in every 1000, uncommon)
 - 4.3.8. Hernia at site of entry
 - 4.3.9. Failure, resulting in unplanned pregnancy: the lifetime failure rate is 1 in 200 (uncommon)
 - 4.3.10. The possibility of a future pregnancy occurring in the fallopian tube if failure occurs
- 4.4. Frequent:
 - 4.4.1. Shoulder-tip pain
 - 4.4.2. Frequency of micturition, dysuria and urinary tract infection





Gleneagles Hospital
HONG KONG
港怡醫院

